

CLAIMS

What is claimed is:

- 5 1. A method of treating a respiratory distress syndrome in a mammal, comprising administering a therapeutically effect amount of an agent that activates surfactant secretion in said mammal.
- 10 2. The method of **Claim 1**, wherein said therapeutic agent is at least one intracellular calcium chelator.
3. The method of **Claim 2**, wherein said intracellular calcium chelator is BAPTA-AM.
- 15 4. The method of **Claim 3**, wherein BAPTA-AM is between 25 and 100 μM .
5. The method of **Claim 1**, wherein said agent comprises inducing an enhanced secretion of surfactant from type II pneumocytes.
- 20 6. The method of **Claim 1**, wherein said agent acts by altering an endoplasmic reticulum free calcium concentration ($[\text{Ca}^{+2}]_i$) in type II pneumocytes.
7. The method of **Claim 1**, wherein said agent is administered by an aerosol, nebulization or liquid instillation.
- 25 8. A method of inhibiting a respiratory distress syndrome in a mammal, comprising administering a therapeutically effect amount of an agent that activates surfactant secretion in said mammal.
- 30 9. The method of **Claim 8**, wherein said agent comprises at least one intracellular calcium chelator.

10. The method of **Claim 9**, wherein said intracellular calcium chelator is BAPTA-AM.

5

11. The method of **Claim 10**, wherein BAPTA-AM is between 25 and 100 μM .

12. The method of **Claim 8**, wherein said agent comprises inducing an enhanced secretion of surfactant from type II pneumocytes.

10

13. The method of **Claim 12**, wherein said agent acts by altering an endoplasmic reticulum free calcium concentration ($[\text{Ca}^{+2}]_i$) in type II pneumocytes.

14. The method of **Claim 8**, wherein said agent is administered by an aerosol, nebulization or liquid instillation.

15